

Buttercups

Trevor D. Israel, Extension Assistant
G. Neil Rhodes, Jr., Professor and Extension Weed Management Specialist
Department of Plant Sciences

Hairy Buttercup *Ranunculus sardous* Crantz

Bulbous Buttercup *Ranunculus bulbosus* L.

Classification and Description

Several species of buttercup are found in Tennessee. Two of the most common are hairy buttercup and bulbous buttercup. They are not native to the United States and are members of the buttercup family (Ranunculaceae). They can be found in pastures, hay fields, roadsides and marginal areas. Although bulbous buttercup is a perennial and hairy buttercup is an annual, they are similar in appearance (Fig. 1). Both species grow from basal rosettes that form in late autumn and winter. Leaves are palmately 3-lobed, 0.75 to 2 inches long and wide, alternate upward, and hairy. Stems are erect, hairy, single or branching from base, and can reach heights of 24 inches. Solitary flowers open in the spring and are pale to bright sulfur yellow, 0.5 to 1 inches wide, with 5 petals and numerous stamens (Fig. 2). The fruit is an aggregate of 10 to 40 achenes in a globose or oblong head, turning brown at maturity. The best way to differentiate the two species is to dig up and examine the underground plant parts. Bulbous buttercup has slightly thickened roots that arise from a corm or “bulb-like” base while hairy buttercup has fibrous roots and does not have a swollen base (Fig. 3). Both species spread by seed but bulbous buttercup can also regenerate from corms.

Problems in Pastures and Hay Fields

Buttercups are not as palatable as desirable forages in a pasture, and cattle will selectively graze around them. However, buttercup species contain the oil protoanemonin, which can cause oral and gastrointestinal irritation. Toxicity varies with species, age, and habitat, but in general, the leaves and stems of flowering plants have the highest concentration. Poisoning is uncommon, but can occur when desirable forages are in short supply. The oil volatilizes rapidly, so plants found in dried hay do not pose toxicity problems. Buttercups can also compete with grasses and increase in density over time. Since buttercups mature in late spring, they can significantly impact the first hay cutting. If not managed, buttercup seed will accumulate in the soil and provide new generations of weeds every year.



Fig. 1. Hairy buttercup (L) and bulbous buttercup (R).



Fig. 2. Buttercup flowers.



Fig. 3. Hairy buttercup (L) is not swollen at the base, whereas bulbous buttercup (R) has an enlarged corm.

Management in Pastures and Hay Fields

Prevention is a crucial component in the management of pasture and hay field weeds. Even just a few mature buttercups growing in a field or brought in from hay bales can provide seed for hundreds of plants the following year. In severe infestations, herbicides are usually needed as part of a control program. Timing is critical for effective buttercup management. Waiting to spray a field when it is a “sea of yellow” is too late because many of the plants have already gone to seed. The best time to spray is in the fall (late October to early December) or late winter to early spring (March to early April). One advantage of spraying in the fall may be time availability. Springtime is very busy for many producers who are in the middle of calving or preparing row crop ground. Another advantage of fall application is there may be fewer gardens and sensitive crops growing that can be injured by herbicide drift. Regardless of a fall or spring application, it is important to have three days where temperatures reach 60 F, so that herbicides can be most effective. Another factor to consider is the height of surrounding forages and thatch. Since buttercup rosettes are low growing, the grass should not be so tall that it prevents the spray from reaching the weeds.

University of Tennessee research indicates that 2,4-D ester applied pre-bloom will adequately control hairy buttercup. However, products containing aminopyralid (GrazonNext HL) or dicamba (Banvel, Clarity, Oracle, others) will give more complete control of bulbous buttercup. As is the case with all pesticide applications, be sure to thoroughly read and follow the label directions. Also, remember that practicing good herbicide stewardship is everyone’s responsibility. For more information regarding herbicide stewardship and reducing off-target damage to sensitive crops, please visit our website: herbicidestewardship.utk.edu.

References

- Everest, J. W., T. A. Powe, Jr., and J. D. Freeman. 1996. pp. 12-13 *in* Poisonous Plants of the Southeastern United States. Alabama Cooperative Extension Service Publ. ANR-975. 51pp.
- Bryson, C. T. and M. S. DeFelice, eds. 2009. p. 268 *in* Weeds of the South, Univ. of Georgia Press, Athens, GA 30602. 468 pp.
- Small, J. K. 1933. p. 522 *in* Manual of the Southeastern Flora, Science Press Printing Co., Lancaster, PA. 1554 pp.

Picture Credits

Fig. 2. Buttercup flowers. Will Phillips, Former Graduate Research Assistant, Plant Sciences Department.

Disclaimer

This publication contains pesticide recommendations that are subject to change at any time. The recommendations in this publication are provided only as a guide. It is always the pesticide applicator's responsibility, by law, to read and follow all current label directions for the specific pesticide being used. The label always takes precedence over the recommendations found in this publication.

Use of trade or brand names in this publication is for clarity and information; it does not imply approval of the product to the exclusion of others that may be of similar, suitable composition, nor does it guarantee or warrant the standard of the product. The author(s), the University of Tennessee Institute of Agriculture and University of Tennessee Extension assume no liability resulting from the use of these recommendations.



AG.TENNESSEE.EDU
Real. Life. Solutions.